

4-DIMENSIONAL TOPOLOGICALLY REGULAR ALEXANDROV SPACES WITH POSITIVE OR NONNEGATIVE CURVATURE AND TORUS SYMMETRY

FERNANDO GALAZ-GARCIA*

ABSTRACT. We point out that a topologically regular Alexandrov space of dimension 4 (with curvature bounded below) and an effective, isometric action of the circle or the 2-torus is locally smooth. This observation implies that the topological and equivariant classifications of compact, simply connected Riemannian 4-manifolds with positive or nonnegative sectional curvature and an effective isometric action of a circle or a 2-torus also hold if we consider topologically regular Alexandrov spaces instead of Riemannian manifolds.

1. INTRODUCTION AND MAIN RESULTS

The interaction between (sectional) curvature and isometric actions of compact Lie groups on Riemannian manifolds plays an important role in the study of compact Riemannian manifolds with positive or nonnegative curvature. Alexandrov spaces (with curvature bounded below) arise naturally in this context as orbit spaces of isometric actions of compact Lie groups on Riemannian manifolds with a lower curvature bound. Alexandrov spaces also occur as Gromov-Hausdorff limits of sequences of Riemannian manifolds. Furthermore, being synthetic generalizations of Riemannian manifolds with curvature bounded below, Alexandrov spaces are geometric objects of intrinsic interest.

It is well-known that the isometry group of a (compact) Riemannian manifold is a (compact) Lie group. Since the same conclusion is true for Alexandrov spaces (cf. [12]), it is of interest to determine which results in equivariant Riemannian geometry have Alexandrov analogues. The simplest Alexandrov spaces for which one may consider this question are those in which every space of directions is homeomorphic to a sphere; following Ge [18], we will say that these spaces are *topologically regular* and shall refer to them as *TR Alexandrov spaces*. These spaces arise, for example, as limits of sequences of n -dimensional Riemannian manifolds with the same lower curvature bound (cf. [24]). Nevertheless, it is not known whether any TR Alexandrov space can be realized as a limit of Riemannian manifolds with the same dimension and lower curvature bound.

Every TR Alexandrov space is isometric to an *Alexandrov manifold*, i.e. a topological manifold equipped with an Alexandrov metric. The converse, however,

Date: August 16, 2012.

2000 *Mathematics Subject Classification.* 53C20.

* The author is part of SFB 878: *Groups, Geometry & Actions*, at the University of Münster.

is not true: The double spherical suspension $\Sigma^2(P^3)$ of the Poincaré homology 3-sphere (P^3, g_0) , with its standard Riemannian metric g_0 of constant positive curvature, is a positively curved Alexandrov manifold homeomorphic to the 5-sphere and is not topologically regular. Ge [18] has classified up to homeomorphism open TR Alexandrov spaces of dimension 4 and nonnegative curvature.

Motivated by the preceding discussion, in this note we show that the topological and equivariant classification of compact, simply connected Riemannian 4-manifolds with positive or nonnegative curvature and an effective isometric action of a circle or a 2-torus (cf. [13, 15, 16, 21, 23, 28, 37]) still holds if instead of considering Riemannian manifolds we consider TR Alexandrov spaces. Before stating our results, recall that a *biquotient* is a quotient of a Lie group G by the two-sided, free action of a subgroup $H \subset G \times G$. If G is equipped with a bi-invariant metric, then the action of H is by isometries and the quotient $G//H$ equipped with the induced metric (of nonnegative curvature) is called a *normal biquotient*.

Theorem A. *Let X^4 be a compact, simply connected, topologically regular Alexandrov space of dimension 4, with an effective, isometric T^2 action.*

- (1) *If $\text{curv}(X^4) \geq 1$, then X^4 is equivariantly homeomorphic to $\mathbb{S}^4$ or $\mathbb{C}P^2$ with a linear action.*
- (2) *If $\text{curv}(X^4) \geq 0$, then X^4 is equivariantly homeomorphic to $\mathbb{S}^2$ or $\mathbb{C}P^2$ with a linear action or to $\mathbb{S}^2 \times \mathbb{S}^2$ or $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ with an action induced by a normal biquotient Riemannian metric.*

Theorem B. *Let X^4 be a compact, simply connected, topologically regular Alexandrov space of dimension 4, with an effective, isometric S^1 action.*

- (1) *If $\text{curv}(X^4) \geq 1$, then X^4 is equivariantly homeomorphic to $\mathbb{S}^4$ or $\mathbb{C}P^2$ with a linear action.*
- (2) *If $\text{curv}(X^4) \geq 0$, then X^4 is equivariantly homeomorphic to $\mathbb{S}^2$ or $\mathbb{C}P^2$ with a linear action or to $\mathbb{S}^2 \times \mathbb{S}^2$ or $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ with an action induced by a normal biquotient Riemannian metric.*

We point out that T^2 is the largest torus that can act isometrically on a simply connected 4-dimensional Alexandrov space (cf. [17]).

Theorems A and B follow from the observation that the torus and circle actions under consideration are locally smooth (see subsection 2.2). This allows us to use Orlik and Raymond's work on compact, simply connected, topological 4-manifolds with locally smooth 2-torus actions (cf. [30]) and Fintushel and Pao's work on compact, simply connected, topological 4-manifolds with locally smooth circle actions (cf. [9, 10, 31]), essentially reducing the proof to the smooth case. The metric and topological tools used in the proofs in the Riemannian case, among them crucial work of Grove and Wilking on extremal knots in an Alexandrov 3-sphere [21], yield the results.

Our note is divided as follows: In Section 2 we collect some background material on isometric actions on Alexandrov spaces and locally smooth actions; in

particular, we observe that the spaces in Theorems A and B are locally smooth. Sections 3 and 4 contain, respectively, the proofs of Theorems A and B.

Acknowledgements. I would like to thank B. Wilking for helpful conversations, and J. Harvey and C. Searle for sharing their preprint [22].

2. PRELIMINARIES

In this section we recall some background material on isometric compact Lie group actions on Alexandrov spaces (cf. [14, 17, 22]) and show that actions of the type considered in Theorems A and B are locally smooth. We assume that the reader is familiar with the basic theory of compact transformation groups (cf. [1]) and of Alexandrov spaces (cf. [3, 4]).

2.1. Equivariant Alexandrov geometry. Let $G \times X \longrightarrow X$, $x \mapsto g(x)$, be a topological action of a topological group G on a Hausdorff topological space X , i.e. let X be a G -space. We will denote the orbit of a point $x \in X$ under the action of G by $G(x) \simeq G/G_x$, where $G_x = \{g \in G : g(x) = x\}$ is the isotropy subgroup of G at x . We will denote the fixed point set of the action by $\text{Fix}(X, G)$. Given a subset $A \subset X$, we will denote its projection under the orbit map $\pi : X \rightarrow X/G$ by A^* . Following this convention, we will denote the orbit space X/G by X^* . We will henceforth assume all actions to be effective. We will say that two G -spaces are *equivalent* if they are equivariantly homeomorphic.

Let (X, dist) be a (finite dimensional) Alexandrov space (with curvature bounded below) with an isometric action of a compact Lie group G . We will denote the space of directions of X at x by $\Sigma_x X$. Given $A \subset \Sigma_x X$, the set of *normal directions* to A , denoted by $\nu(A)$, is defined by

$$\nu(A) = \{v \in \Sigma_x X : \text{dist}(v, w) = \text{diam}(\Sigma_x X)/2 \text{ for all } w \in A\}.$$

Let $S_x \subset \Sigma_x X$ be the unit tangent space to the orbit $G(x)$ at x and suppose that $\dim(G(x)) > 0$. The set $v_x = \nu(S_x)$ is a compact, totally geodesic Alexandrov subspace of $\Sigma_x X$ with curvature bounded below by 1, and the space of directions $\Sigma_x X$ is isometric to the join $S_x * v_x$ with the standard join metric. Moreover, either v_x is connected or it contains exactly two points at distance π (cf. [17]).

The slice and principal orbit theorems, as well as Kleiner's isotropy lemma for isometric actions of compact Lie groups on Riemannian manifolds (cf. [19]) also hold for Alexandrov spaces (cf. [14, 22]). In particular, if a compact Lie group G acts by isometries on an Alexandrov space X , then a slice at x is G_x -equivariantly homeomorphic to the cone on v_x , the space of normal directions to the orbit. In other words, for any $x \in X$ there is some $r_0 > 0$ such that for all $r < r_0$ there is an equivariant homeomorphism $\varphi : G \times_{G_x} K(v_x) \rightarrow B_r(G(x))$ (cf. [22]). It follows that $\Sigma_{x^*} X^*$, the space of directions of the orbit space X^* at x^* , is isometric to v_x/G_x .

2.2. Locally smooth actions. Let G be a compact Lie group and let X be a G -space. Let $G(x)$ be an orbit of type G/H , for some $x \in X$. Recall that a *tube* about

$G(x)$ is a G -equivariant embedding (homeomorphism into)

$$\varphi : G \times_H A \longrightarrow X,$$

onto an open neighborhood of $G(X)$ in X , where A is some space on which H acts. If A is homeomorphic to Euclidean space V and H acts orthogonally on V , then we say that the tube

$$\varphi : G \times_H V \longrightarrow X$$

is a *linear tube*. A slice S at $x \in X$ is called a *linear slice* if the canonically associated tube

$$\begin{aligned} G \times_{G_x} S &\longrightarrow X \\ [g, s] &\mapsto g(s) \end{aligned}$$

is equivalent to a linear tube, i.e. if the G_x -space S is equivalent to an orthogonal G_x -space. We say that a G -space is *locally smooth* if there exists a linear tube about each orbit. Since $G \times_H V$ is a V -bundle over G/H , it's a topological manifold. Thus, any locally smooth G -space must be a topological manifold. If the G -space X is locally smooth and if x is a fixed-point, then a neighborhood of x in X is equivalent to an orthogonal action. Hence, in this situation, the fixed point set of the G action is a topological submanifold of X . There exist examples of topological actions on manifolds which are not locally smooth (cf. [1, Ch. IV]). We now show that the G -spaces considered in Theorems A and B are locally smooth.

Lemma 2.1. *If X is a TR Alexandrov space of dimension 4 with an effective, isometric action of T^2 or S^1 , then X is locally smooth.*

Proof. Let X be a TR Alexandrov space with an effective, isometric action of G , with G isomorphic to T^2 or S^1 . By the slice theorem, it suffices to show that the isotropy action on the normal space of directions to any orbit is equivalent to a linear action on a sphere. Suppose first that $G \cong T^2$. In this case, the only possible non-principal isotropy subgroups of the torus action are conjugate to S^1 or T^2 (cf. [30]). The space of directions normal to the orbit is, respectively, homeomorphic to $\mathbb{S}^2$ or to $\mathbb{S}^3$ and it is well-known that topological S^1 actions on $\mathbb{S}^2$ and topological T^2 actions on $\mathbb{S}^3$ are equivalent to linear actions.

Suppose now that $G \cong S^1$. In this case, the only possible non-principal isotropy subgroups are finite cyclic groups $\mathbb{Z}_k$, $k \geq 2$, or S^1 . The space of directions normal to the orbit is, respectively, homeomorphic to $\mathbb{S}^2$ or to $\mathbb{S}^3$. In the first case, $\mathbb{Z}_k$ acts by homeomorphisms on $\mathbb{S}^2$. By work of de Kerékjártó [7], Brouwer [2] and Eilenberg [8], this action is equivalent to an orthogonal action (cf. [6]). In the second case, S^1 acts by homeomorphisms on $\mathbb{S}^3$ and, by work of Raymond [36, Theorem 6], this action is equivalent to an orthogonal action. $\square$

Recall that if M is a smooth compact manifold with a smooth S^1 action, then $\chi(\text{Fix}(M, S^1))$, the Euler characteristic of the fixed point set, is the Euler characteristic of M (cf. [29, Theorem 5.5-(1)]). An inspection of Kobayashi's proof shows that in our case, where we consider locally smooth circle actions on compact, topological manifolds, the same conclusion holds.

Proposition 2.2. *If $S^1 \times X \longrightarrow X$ is a locally smooth circle action on a compact topological manifold X , then*

$$\chi(\text{Fix}(X, S^1)) = \chi(X).$$

3. PROOF OF THEOREM A

By work of Orlik and Raymond [30], X is homeomorphic to a connected sum of copies of $\mathbb{S}^2 \times \mathbb{S}^2$, $\pm \mathbb{C}P^2$ and $\mathbb{S}^4$. Moreover, the T^2 action is equivalent to a smooth T^2 action on X , where X has the standard smooth structure induced via connected sum by the standard smooth structure on each one of $\mathbb{S}^2 \times \mathbb{S}^2$, $\mathbb{C}P^2$ and $\mathbb{S}^4$. The orbit space X^* is homeomorphic to a 2-disk and the action has at least 2 fixed points. On the other hand, proceeding as in [16], we conclude that the action has at most 3 fixed points, if $\text{curv}(X) \geq 1$, and at most 4 fixed points, if $\text{curv}(X) \geq 0$. It follows from [30] that, if the action has 2 or 3 fixed points, then X is equivariantly homeomorphic, respectively, to $\mathbb{S}^4$ or $\mathbb{C}P^2$ equipped with a linear action; if the action has 4 fixed points, then X is equivariantly homeomorphic to $\mathbb{S}^2 \times \mathbb{S}^2$ or $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ with a smooth T^2 action. By [15], any smooth T^2 action on $\mathbb{S}^2 \times \mathbb{S}^2$ or $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ is equivariantly diffeomorphic to an isometric action on a normal biquotient.

4. PROOF OF THEOREM B

By work of Fintushel [9, 10], Pao [31] and Perelman [33, 34, 26, 27], X is homeomorphic to a connected sum of copies of $\mathbb{S}^2 \times \mathbb{S}^2$, $\pm \mathbb{C}P^2$ and $\mathbb{S}^4$. Moreover, the S^1 action is equivalent to a smooth S^1 action on X equipped with the standard smooth structure induced via connected sum by the standard smooth structure on each one of $\mathbb{S}^2 \times \mathbb{S}^2$, $\mathbb{C}P^2$ and $\mathbb{S}^4$. The action is determined by Fintushel's orbit space weights (cf. Theorem 13.2 in [10]). Observe that $\chi(\text{Fix}(X, S^1)) = \chi(X)$ via a well-known theorem of Kobayashi for smooth circle actions on compact smooth manifolds (cf. also Proposition 2.2).

It is well-known that the only possible orbit types are principal orbits, fixed points and exceptional orbits. For convenience, in this section we will also denote the set of fixed points by F ; the set of exceptional orbits will be denoted by E . The orbit space X^* is a simply connected topological 3-manifold with $\partial X^* \subset F^*$, the set $F^* - \partial X^*$ of isolated fixed points is finite and F^* is nonempty. The components of ∂X^* are 2-spheres and the closure $\overline{E}^*$ of E^* is a collection of polyhedral arcs and simple closed curves in X^* . The components of E^* are open arcs on which orbit types are constant, and these arcs have closures with distinct endpoints in $F^* - \partial X^*$, so that $\overline{E}^* \subset E^* \cup F^*$. It follows from [9, Theorem 7.1] and the validity of the Poincaré conjecture that, if $\overline{E}^*$ contains no simple closed curves, then the S^1 action on X extends to a locally smooth action of T^2 . If $\overline{E}^*$ contains a simple closed curve K^* , then the S^1 action extends to a locally smooth T^2 action if and only if $E^* \cup F^* = K^*$ and K^* is unknotted in $X^* \simeq \mathbb{S}^3$. Observe that the simple closed curves in $\overline{E}^*$ are tame knots in X^* .

Proof of part (1) - Positive curvature. Let X be a compact, simply connected TR Alexandrov space with $\text{curv} \geq 1$ and an isometric circle action.

We derive first the topological classification. By Freedman [11], it suffices to show that $2 \leq \chi(X) \leq 3$. The lower bound on $\chi(X)$ follows from Poincaré duality. We obtain the upper bound on $\chi(X)$ by showing that the fixed point set of the action has Euler characteristic at most 3. Suppose that the fixed point set F contains a 2-sphere, which must be a boundary component of the orbit space X^* . Since X^* is a positively curved Alexandrov space, the distance function to ∂X^* is strictly concave and it follows from Perelman's Soul Theorem for Alexandrov spaces [32] that the set at maximal distance from ∂X^* is a point x_0^* . The distance function to ∂X^* has no critical points in $X^* - \{\partial X^* \cup \{p_0^*\}\}$ and, proceeding as in [20, Lemma 1.1-(iii)], one concludes that the points in $X^* - \{\partial X^* \cup \{p_0^*\}\}$ correspond to orbits with trivial isotropy. Hence, the fixed point set of the action is either a 2-sphere, or a 2-sphere and an isolated fixed point. Suppose now that the fixed point set consists only of isolated fixed points. A standard triangle comparison argument shows that there can be at most three such points. Therefore, $\chi(F) \leq 3$.

We now derive the equivariant classification. By work of Orlik and Raymond [30], any topological T^2 action on $\mathbb{S}^4$ or $\mathbb{C}P^2$ is equivalent to a linear action. Thus, to obtain the desired conclusion, it suffices to show that the S^1 action extends to a T^2 action. By the discussion at the beginning of this section, this is immediate when the fixed point set contains a 2-sphere, since in this case the set E of exceptional orbits is empty.

If F consists of only isolated points and $\overline{E}^*$ contains a simple closed curve K^* , we must verify that $K^* = E^* \cup F^*$ and K^* is unknotted in $X^* \simeq \mathbb{S}^3$. To do so, we may proceed along the lines of [21]; for the sake of completeness, we briefly sketch here the relevant ideas (details can be found in [21]). One first verifies that the canonical two-fold branched cover $X_2^*(K^*)$ of X^* , with branching set K^* , is an Alexandrov space with the same lower curvature bound as the orbit space X^* . To see this, let $C^* \subset K^*$ be the closure of the open arcs in K^* whose isotropy is not $\mathbb{Z}_2$. Observe that $X_2^*(K^*)$ is an Alexandrov space in a neighborhood of $K^* - C^*$, since there it is locally isometric to the quotient of a neighborhood of an orbit of the S^1 action on X . On the other hand, since the isotropy group is constant in the interior of minimal geodesics joining any two orbits, the set $X^* - C^*$ is convex, and so is the set $X_2^*(K^*) - C^*$. Since any geodesic triangle in $X_2^*(K^*)$ is the limit of geodesic triangles in $X_2^*(K^*) - C^*$, the conclusion follows. One may then use the weighted orbit space restrictions in [9] and the fact, due to Grove and Wilking [21], that $\pi_1(X_2^*(K^*)) = 0$ if K^* is unknotted, and $|\pi_1(X_2^*(K^*))| \geq 3$ otherwise, to conclude that $K^* = E^* \cup F^*$ and K^* is unknotted in $X^* \simeq \mathbb{S}^3$. Therefore, as recalled at the beginning of this section, the S^1 action can be extended to a locally smooth T^2 action.

Proof of part (2) - Nonnegative curvature. Let X be a compact TR Alexandrov space of dimension 4 with $\text{curv}(X) \geq 0$ and an isometric circle action.

As in the positively curved case, to derive the topological classification it suffices to show that $2 \leq \chi(X) \leq 4$, with the lower bound immediately following

from Poincaré duality. We have three cases to consider: when the fixed point set consists only of isolated points, when there are 2-dimensional components and isolated fixed points, and where there are only 2-dimensional components. In the first case, a comparison argument shows that there can be at most four such points. In the second case, the metric argument used in [37] implies that there can only be at most two isolated fixed points and one 2-dimensional component. In the third case, an analog of the Searle-Yang Double Soul Theorem in [37] holds in the case at hand and implies that if there are at least two 2-dimensional fixed point set components, there must be exactly two and there cannot be isolated fixed points:

Theorem 4.1. *Let X be a complete TR regular Alexandrov space of nonnegative curvature equipped with an isometric, locally smooth S^1 action. Let F be the fixed point set of the S^1 action on X . Suppose that F contains two connected codimension 2 components Y and Z with one of them being compact. Then Y is isometric to Z , $F = Y \cup Z$ and X is homeomorphic to an $\mathbb{S}^2$ -bundle over Y .*

The proof of this theorem follows from observing that X^* is a TR Alexandrov space of nonnegative curvature and, by the analysis in [38] (cf. also [5]), X^* must split isometrically as $Y \times I$. In particular, in the 4-dimensional case, the orbit space must be isometric to $\mathbb{S}^2 \times I$.

Therefore, the fixed point set F of the action can only consist of four isolated fixed points, a 2-sphere and at most two isolated fixed points, or two 2-spheres. In all cases $\chi(F) \leq 4$.

Let X be isometric to one of $\mathbb{S}^4$, $\mathbb{C}P^2$ or $\mathbb{S}^2 \times \mathbb{S}^2$ or $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ equipped with an Alexandrov metric of nonnegative curvature and an isometric circle action. Recall that, by work of Orlik and Raymond [30], any locally smooth smooth T^2 action on $\mathbb{S}^4$, $\mathbb{C}P^2$ or $\mathbb{S}^2 \times \mathbb{S}^2$ or $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ is equivalent to a smooth action. By [30], any smooth T^2 action is equivalent to a linear action and, by [15], any smooth T^2 action on $\mathbb{S}^2 \times \mathbb{S}^2$ or $\mathbb{C}P^2 \# \pm \mathbb{C}P^2$ is equivalent to an isometric action on a normal biquotient. Thus, to obtain the equivariant classification of locally smooth circle actions on X it suffices, by the discussion at the beginning of this section, to verify that if $K^* \subset \overline{E}^*$ is a simple closed curve, then $K^* = E^* \cup F^*$ and K^* is unknotted in $X^* \simeq \mathbb{S}^3$. This follows as in the positively curved case.

REFERENCES

1. Bredon, G. E., *Introduction to compact transformation groups*, Pure and Applied Mathematics, vol. 46, Academic Press, New York-London, 1972.
2. Brouwer, L.E.J., *Über die periodischen transformationen der Kugel*, Math. Ann. **80** (1919), 39–41.
3. Burago, D., Burago, Y. and Ivanov, S., *A course in metric geometry*, Graduate Studies in Mathematics, vol. 33, American Mathematical Society, Providence, RI, 2001.
4. Burago, Y., Gromov, M. and Perel'man, G., *A. D. Aleksandrov spaces with curvatures bounded below*, (Russian) Uspekhi Mat. Nauk **47** (1992), no. 2(284), 3–51, 222; translation in Russian Math. Surveys **47** (1992), no. 2, 1–58.
5. Cao, J. and Ge, J., *A simple proof of Perelman's collapsing theorem for 3-manifolds*, J. Geom. Anal. **21** (2011), no. 4, 807–86.

6. Constantin, A. and Kolev, B., *The theorem of Kerékjártó on periodic homeomorphisms of the disc and the sphere*, Enseign. Math. (2) **40** (1994), no. 3-4, 193–204.
7. de Kerékjártó, B., *Sur les groupes compacts de transformations topologiques des surfaces*, Acta Math. **74** (1941), 129–173.
8. Eilenberg, S., *Sur les transformations periodiques de la surface de la sphere*, Fund. Math. **22** (1934), 28–44.
9. Fintushel, R., *Circle actions on simply connected 4-manifolds*, Trans. Amer. Math. Soc. **230** (1977), 147–171.
10. Fintushel, R., *Classification of circle actions on 4-manifolds*, Trans. Amer. Math. Soc. **242** (1978), 377–390.
11. Freedman, M., *The topology of four-dimensional manifolds*, J. Differential. Geom. **17** (1982) 357–453.
12. Fukaya, K. and Yamaguchi, T., *Isometry groups of singular spaces*, Math. Z. **216** (1994), no. 1, 31–44.
13. Galaz-Garcia, F., *Nonnegatively curved fixed point homogeneous manifolds in low dimensions*, Geometriae Dedicata, to appear.
14. Galaz-Garcia, F. and Guijarro, L., *Isometry groups of Alexandrov spaces*, preprint (2012), arXiv:1109.4878v2 [math.DG].
15. Galaz-Garcia, F. and Kerin, M., *Cohomogeneity-two torus actions on non-negatively curved manifolds of low dimension*, preprint (2011), arXiv:1111.1640v1 [math.DG].
16. Galaz-Garcia, F. and Searle, C., *Low-dimensional manifolds with non-negative curvature and maximal symmetry rank*, Proc. Amer. Math. Soc., **139** (2011), 2559–2564.
17. Galaz-Garcia, F. and Searle, C., *Cohomogeneity one Alexandrov spaces*, Transform. Groups **16** (2011), no. 1, 91–107.
18. Ge, J., *Soul theorem for 4-dimensional topologically regular open nonnegatively curved Alexandrov spaces* Proc. Amer. Math. Soc. **139** (2011), no. 12, 4435–4443.
19. Grove, K., *Geometry of and via symmetries*, Conformal, Riemannian and Lagrangian geometry (Knoxville, TN, 2000), 31–53, Univ. Lecture Ser., 27, Amer. Math. Soc., Providence, RI, 2002.
20. Grove, K. and Searle, C., *Positively curved manifolds with maximal symmetry-rank*, J. Pure Appl. Algebra **91** (1994), no. 1-3, 137–142.
21. Grove, K. and Wilking, B., *A knot characterization and 1-connected nonnegatively curved 4-manifolds with circle symmetry*, preprint (2012).
22. Harvey, J. and Searle, C., *Isotropy and the slice lemma in Alexandrov spaces*, preprint (2012).
23. Hsiang, W.Y. and Kleiner, B., *On the topology of positively curved 4-manifolds with symmetry*, J. Differential Geom. **29** (1989), no. 3, 615–621.
24. Kapovitch, V., *Regularity of limits of noncollapsing sequences of manifolds*, Geom. Funct. Anal. **12** (2002), no. 1, 121–137.
25. de Kerékjártó, B., *Sur les groupes compacts de transformations topologiques des surfaces*, (French) Acta Math. **74**, (1941). 129–173.
26. Kleiner, B. and Lott, J., *Notes on Perelman’s papers*, Preprint (2006), arXiv:math/0605667v2 [math.DG].
27. Morgan, J. and Tian, G., *Ricci flow and the Poincaré conjecture*, Clay Mathematics Monographs, vol. 3, American Mathematical Society, Providence, RI, 2007.
28. Kleiner, B., *Riemannian four-manifolds with nonnegative curvature and continuous symmetry*, Ph.D. thesis, University of California, Berkeley, 1990.
29. Kobayashi, S., *Transformation groups in differential geometry*, Reprint of the 1972 edition. Classics in Mathematics. Springer-Verlag, Berlin, 1995.
30. Orlik, P. and Raymond, F., *Actions of the torus on 4-manifolds I*, Trans. Amer. Math. Soc. **152** 531–559 (1970).
31. Pao, P. S., *Nonlinear circle actions on the 4-sphere and twisting spun knots*, Topology **17** (1978), no. 3, 291–296.
32. Perelman, G., *Alexandrov’s spaces with curvatures bonded from below II*, Preprint.

- 33. Perelman, G., *The entropy formula for the Ricci flow and its geometric applications*, preprint (2002), arXiv:math/0211159v1 [math.DG].
- 34. Perelman, G., *Ricci flow with surgery on three-manifolds*, preprint (2003), arXiv:math/0303109v1 [math.DG].
- 35. Perelman, G., *Finite extinction time for the solutions to the Ricci flow on certain three-manifolds*, preprint (2003), arXiv:math/0307245v1 [math.DG].
- 36. Raymond, F., *Classification of the actions of the circle on 3-manifolds*, Trans. Amer. Math. Soc. **131** (1968), 51–78.
- 37. Searle, C. and Yang, D., *On the topology of non-negatively curved simply connected 4-manifolds with continuous symmetry*, Duke Math. J. **74** (1994), no. 2, 547–556.
- 38. Shioya, T. and Yamaguchi, T., *Collapsing three-manifolds under a lower curvature bound*, J. Differential Geom. **56** (2000), no. 1, 1–66.

MATHEMATISCHES INSTITUT, WWU MÜNSTER, GERMANY
E-mail address: f.galaz-garcia@uni-muenster.de